




## Human Fibroblast Growth Factor 2 (FGF2) Protein, Recombinant

### I. For sale

Product name	Catalog #	Size
Human Fibroblast Growth Factor 2 (FGF2) Protein, Recombinant	P01F0003	10ug
		50ug
		500ug
		1mg

### II. Product Description

Other Names	BFGF; FGFB; FGF-2; HBGF-2
Protein & NCBI Number	D9ZGF5, NM_001361665.2
Host	E.coli
Express Region	Met1-Ser155
Protein Sequence	MAAGSITLTPALPEDGGSGAFPPGHFKDPKRLYCKNGGFFLRIH PDGRVDGVREKSDPHIKLQLQAEERGVSISIKGVCANRYLAMKEDGRLLASKCVTDECF FFERLESNNYNTYRSRKYTSWYVALKRTGQYKLGSKTGPQGKAILFLPMSAKS
Molecular Weight	The protein consists of 281 amino acids (including the fusion tag), with a predicted molecular weight of 31.5 kDa, which matches the actual molecular weight.
Fusion Tag	6×His-SUMO (N-terminus)
Purity	≥95% SDS-PAGE
Physical Property	Liquid
components	0.01M PBS+20% glycerol, sterile solution.
Storage & Stability	After aliquoting, the stability of the samples can be maintained for up to 6 months at -20°C to -80°C, avoiding repeated freeze-thaw cycles.
Applications	Antibody preparation, immunoassay (ELISA, WB), subcellular localization and interaction protein identification, etc.
Lead Time	5 to 10 business days; 2 to 3 days for stock products
Figure. SDS-PAGE	 <p>Bis-Tris (MOPS) SDS-PAGE</p>



### III. Storage and Transportation

Transport at 2-8°C, product is stable for up to twelve months from date of receipt under sterile conditions at -20°C to -80°C.

### IV. Notes

This product is for research use only. Please wear laboratory attire and disposable gloves when handling.

### V. Background

FGF-2, also known as basic fibroblast growth factor (bFGF), is an important member of fibroblast growth factor (FGF) family. FGF-2 is a cationic polypeptide with a molecular weight of 16 ~ 18000 and an isoelectric point of 9.6. FGF2 can be produced by vascular endothelial cells, retinal pigment epithelial cells, photoreceptor cells, Müller cells and astrocytes. It widely exists in a variety of tissues in vivo and mainly plays a role through autocrine and paracrine. The signal pathway induced by FGF2 is necessary for normal cell growth and differentiation. It exists in almost all cells. FGF2 binds to FGFR, which makes the receptor dimerize and tyrosine kinase is activated, triggering a series of intracellular phosphorylation cascade reactions to regulate cell growth Differentiation and apoptosis. Under normal conditions, FGF-2 binds to heparin and does not produce biological effects. However, in some pathological cases, the integrity of cells is destroyed, which can release the stored form of FGF-2, promote angiogenesis and participate in the process of tissue repair. FGF-2 and FGFR are almost distributed in various tissues of the whole body. FGF-2 is the strongest known cytokine. It plays an important role in promoting angiogenesis, wound healing, tissue injury repair, neuroprotection, embryonic development and tumor formation. FGF-2 has two main functions, inducing endothelial cell germination and proliferation and increasing vascular permeability. In addition, studies have shown that FGF2 is also closely related to depression.

### VI. References

1. 童安莉.成纤维细胞生长因子 2 激活肾上腺皮质癌 H295R 细胞的 ERK、AKT 信号通路.癌症进展,2012,10(01):94-97.
2. 朱华锋,汪春兰,赵宇.VEGF 和 FGF-2 在血管生成中的协同作用研究进展.中华整形外科杂志,2006(01):72-75.
3. 谷中秀,吴绵绵,郭芳,赵少贞,张琰.FGF2 与新生血管性视网膜病变的研究进展.生命科学仪器,2018,16(03):16-22.
4. Germán Andrés et al. A pro - inflammatory signature mediates FGF2 - induced angiogenesis. Journal of Cellular and Molecular Medicine, 2009, 13(8b) : 2083-2108.
5. Barrientos S,Brem H,Stojadinovic O et al. Clinical application of growth factors and cytokines in



---

wound healing. *Wound Repair and Regeneration*, 2014, 22(5) : 569-578.

6. Murakami Shinya. [Periodontal regeneration by FGF2]. *Clinical calcium*, 2007, 17(2):249-55.

